

z -Scaling and Prompt J/ψ Production at High- p_T in $\bar{p}p$ at Tevatron

M.V. Tokarev*

Joint Institute for Nuclear Research, Dubna, Moscow region, 141980, Russia

New data on high- p_T spectra of J/ψ measured by the CDF Collaboration at the Tevatron are analyzed in the framework of z -scaling. F -dependence of z -presentation of the data is used to construct the scaling function of J/ψ . The energy independence of the scaling function is used to predict inclusive cross section of prompt J/ψ production in $\bar{p}p$ collisions at high- p_T . The concept of z -scaling is considered to reflect the general features of high- p_T particle production in hadron-hadron and hadron-nucleus collisions. Violation of z -scaling is suggested to consider as a signature of new physics.

1. INTRODUCTION

Primary goal of all new accelerators is to discover new physics phenomena. They can give impulse for development of physics theory and extend our understanding of micro world. The origin of particle structure at small scales, distributions of mass, charge and spin in space-time, features of particle-to-particle transitions, are of the most enigmatic and least understandable physics phenomena. Therefore, search for scaling regularities in high energy particle collisions is a subject of intense investigations^{1,2,3,4,5,6,7,8,9,10}.

Properties of particle formation are believed to reveal themselves more clearly at high energy $\sqrt{s}$ and transverse momenta p_T . It is considered also that partons produced in hard scattering retain information about primary collision during hadronization. It is known that the mechanism of particle formation is modified by nuclear medium and can be sensitive to the phase transition of nuclear matter. Therefore, the features of high- p_T single inclusive particle spectra of hadron-hadron and hadron-nucleus collisions are of interest to search for new physics phenomena in elementary processes (quark compositeness, fractal space-time, extra dimensions), signatures of exotic state of nuclear matter (phase transitions, quark-gluon plasma), complementary restrictions for theory (nonperturbative QCD effects, physics beyond Standard Model).

New feature (z -scaling) of high- p_T particle production in hadron-hadron and hadron-nucleus collisions at high energies was established in^{11,12}. The scaling function ψ and scaling variable z are expressed via experimental quantities such as the inclusive cross section $E d^3\sigma/dp^3$ and the multiplicity density of charged particles $dN/d\eta$. The z -presentation of data is found to reveal symmetry properties (energy independence, A- and F-dependence, power law). The function $\psi(z)$ is interpreted as the probability density to produce a particle with a formation length z . The concept of z -scaling and the method of data analysis are developed for description of different particles (charged and neutral hadrons, direct photons jets) produced in high energy hadron-hadron and hadron-nucleus interactions. The proposed method is complementary to a method of direct calculations developed in the framework of QCD and methods based on the Monte Carlo generators. Therefore we consider that the use of the method allow to reduce some theoretical uncertainties which are ambiguously estimated by theory.

Recently new data¹⁴ on cross section of prompt J/ψ production in $\bar{p}p$ collisions in the central rapidity range $|\eta_{J/\psi}| < 0.6$ at $\sqrt{s} = 1.96$ TeV are measured by the CDF Collaboration. The data and the other one¹³ obtained in Run I give new possibility to verify models of J/ψ production mechanisms. In the color singlet model¹⁵ the charmonium meson retains the quantum numbers of the produced $\bar{c}c$ pair and each J/ψ state is directly produced in hard scattering of color singlet via the corresponding subprocess. As shown in¹³ the model predictions are in a disagreement with experimental data on J/ψ and $\psi(2S)$ production rates. A color octet model^{16,17} was suggested to explain the discrepancy. The color octet mechanism takes into account the production of $\bar{c}c$ pair in a color octet configuration. The color octet state evolves into a color singlet state via emission of a soft gluon. The color octet contributions were found in¹⁸ from a fit to CDF data¹³.

We use a complementary method of data analysis and present the results of analysis of CDF data¹⁴ on prompt J/ψ production in $\bar{p}p$ collisions in Run I and Run II at Tevatron in the framework of z -scaling. Properties of z -scaling, the energy independence and F -dependence, are used to construct the scaling function for J/ψ . The inclusive cross sections of J/ψ production in $\bar{p}p$ collisions at high- p_T are estimated. The results can be useful to test phenomenological models of J/ψ production mechanism and obtain new restrictions on a gluon distribution function. The possibility to search for the scaling violation at the Tevatron is suggested as well.

* e-mail: tokarev@sunhe.jinr.ru

2. Z-SCALING

In the section underlying ideas of z -scaling, a general scheme of data z -presentation and physical meaning of the scaling function $\psi(z)$ and the scaling variable z are discussed.

2.1. Basic principles: locality, self-similarity, fractality

The idea of z -scaling is based on the assumptions⁶ that gross feature of inclusive particle distribution of the process (1) at high energies can be described in terms of the corresponding kinematic characteristics

$$M_1 + M_2 \rightarrow m_1 + X \quad (1)$$

of the constituent subprocess written in the symbolic form (2)

$$(x_1 M_1) + (x_2 M_2) \rightarrow m_1 + (x_1 M_1 + x_2 M_2 + m_2) \quad (2)$$

satisfying the condition

$$(x_1 P_1 + x_2 P_2 - p)^2 = (x_1 M_1 + x_2 M_2 + m_2)^2. \quad (3)$$

The equation is the expression of locality of hadron interaction at constituent level. The x_1 and x_2 are fractions of the incoming momenta P_1 and P_2 of the colliding objects with the masses M_1 and M_2 . They determine the minimum energy, which is necessary for production of the secondary particle with the mass m_1 and the four-momentum p . The parameter m_2 is introduced to satisfy the internal conservation laws (for baryon number, isospin, strangeness, and so on).

The equation (3) reflects minimum recoil mass hypothesis in the elementary subprocess. To connect kinematic and structural characteristics of the interaction, the quantity Ω is introduced. It is chosen in the form

$$\Omega(x_1, x_2) = m(1 - x_1)^{\delta_1}(1 - x_2)^{\delta_2}, \quad (4)$$

where m is a mass constant and δ_1 and δ_2 are factors relating to the anomalous fractal dimensions of the colliding objects. The fractions x_1 and x_2 are determined to maximize the value of $\Omega(x_1, x_2)$, simultaneously fulfilling the condition (3)

$$d\Omega(x_1, x_2)/dx_1|_{x_2=x_2(x_1)} = 0. \quad (5)$$

The fractions x_1 and x_2 are equal to unity along the phase space limit and cover the full phase space accessible at any energy.

Self-similarity is a scale-invariant property connected with dropping of certain dimensional quantities out of physical picture of the interactions. It means that dimensionless quantities for the description of physical processes are used. The scaling function $\psi(z)$ depends in a self-similar manner on the single dimensionless variable z . It is expressed via the invariant cross section $E d^3\sigma/dp^3$ as follows

$$\psi(z) = -\frac{\pi s}{(dN/d\eta)\sigma_{in}} J^{-1} E \frac{d^3\sigma}{dp^3} \quad (6)$$

Here, s is the center-of-mass collision energy squared, σ_{in} is the inelastic cross section, J is the corresponding Jacobian. The factor J is the known function of the kinematic variables, the momenta and masses of the colliding and produced particles.

The function $\psi(z)$ is normalized as follows

$$\int_0^\infty \psi(z) dz = 1. \quad (7)$$

The relation allows us to interpret the function $\psi(z)$ as a probability density to produce a particle with the corresponding value of the variable z . We would like to emphasize that the existence of the function $\psi(z)$ depending on a single dimensionless variable z and revealing scaling properties is not evident in advance. Therefore the method proposed for data analysis in z -presentation could be only proved a posteriori.

Principle of fractality states that variables used in the description of the process diverge in terms of the resolution. This property is characteristic for the scaling variable

$$z = z_0 \Omega^{-1}, \quad (8)$$

where

$$z_0 = \sqrt{\hat{s}_\perp}/(dN/d\eta). \quad (9)$$

The variable z has character of a fractal measure. For the given production process (1), its finite part z_0 is the ratio of the transverse energy released in the binary collision of constituents (2) and the average multiplicity density $dN/d\eta|_{\eta=0}$. The divergent part Ω^{-1} describes the resolution at which the collision of the constituents can be singled out of this process. The $\Omega(x_1, x_2)$ represents relative number of all initial configurations containing the constituents which carry fractions x_1 and x_2 of the incoming momenta. The δ_1 and δ_2 are the anomalous fractal dimensions of the colliding objects (hadrons or nuclei). The momentum fractions x_1 and x_2 are determined in a way to minimize the resolution $\Omega^{-1}(x_1, x_2)$ of the fractal measure z with respect to all possible sub-processes (2) subjected to the condition (3). The variable z was interpreted as a particle formation length.

The scaling function of high- p_T particle production, as shown below, is described by the power law, $\psi(z) \sim z^{-\beta}$. Both quantities, ψ and z , are scale dependent. Therefore we consider the high energy hadron-hadron interactions as interactions of fractals. In the asymptotic region the internal structure of particles, interactions of their constituents and mechanism of real particle formation manifest self-similarity and fractality over a wide scale range.

2.2. Prompt J/ψ production in $\bar{p}p$

In this section we analyze the data^{13,14} on inclusive cross section of prompt J/ψ production in $\bar{p}p$ collisions. Firstly the experimental data¹⁹ on inclusive cross section of π^0 -mesons produced in $\bar{p}p$ collisions obtained by the UA1 Collaboration at $\sqrt{s} = 540$ GeV are used to construct z -presentation of the data. The scaling function of π^0 is shown in Fig.1(a). The dashed line is the fit of the data. As seen from Fig.1(a) the scaling function is described by the power law, $\psi(z) \sim z^{-\beta}$, on the log-log scale. The value of the slope parameter β is found to be 5.77 ± 0.02 over a wide p_T range. Secondly the F -dependence of z -scaling is used to compare the scaling function of π^0 and J/ψ . We assume that scaling functions of produced particles with different flavor content has the same asymptotic behavior^{11,12}. The transformation of variable z and scaling function ψ in the form

$$z \rightarrow \alpha_F \cdot z, \quad \psi \rightarrow \alpha_F^{-1} \cdot \psi \quad (10)$$

was used. The transformation parameter α_F is found to be 2.25. As seen from Fig.1(a) the scaling function of J/ψ at $z > 10$ reveals the asymptotic behavior such as the similar one for π^0 . We would like to note that the CDF Run I and Run II data demonstrate a good matching in the overlapping region in z -presentation. The power law for the scaling function is observed up to $z \simeq 40$. Therefore the kinematic range of more preferable for search for the scaling violation is found to be for the range $p_T > 30$ GeV/c at $\sqrt{s} = 1960$ GeV.

The energy independence of the constructed scaling function is used to calculate the dependence of inclusive cross section $Ed^3\sigma/dp^3$ of J/ψ production in $\bar{p}p$ collisions on the transverse momentum p_T in the central rapidity range, $\theta \simeq 90^\circ$, at the energy $\sqrt{s} = 63, 200, 630, 1800$ and 1960 GeV. The calculated results are shown by the dashed lines in Fig.1(b). Points ($\star, +$) shown in Fig.1(b) are the corresponding CDF Run I and Run II data. As seen from Fig.1(b) the strong dependence of inclusive cross section on $\sqrt{s}$ increases with p_T .

3. $z - p_T$ PLOT

The scaling function $\psi(z)$ demonstrates the universality over a wide kinematic range. It describes two regimes of particle formation observed at higher energies. The first one reveals exponential behavior at low z and the second one is described by the power law, $\psi(z) \sim z^{-\beta}$, at high z .

Asymptotic behavior of the scaling function is described by the slope parameter β , which depends on the anomalous fractal dimensions $\delta_{1,2}$ of colliding objects. We assume that new physics effects at small scales (high- z) should be characterized by the change of the fractal dimension (" δ -jump") and consequently the other value of the slope parameter β at high z . To determine the kinematic range which is of more preferable for experimental searching for z -scaling violation the $z - p_T$ plot can be used. Figure 2 shows the $z - p_T$ plot, the dependence of the variable z on the transverse momentum p_T of J/ψ -mesons produced in $\bar{p}p$ collisions at $\sqrt{s} = 63 - 14000$ GeV and $\theta_{cms} = 90^\circ$. As seen from Fig.1(a) the asymptotic value of the scaling function $\psi(z)$ at $z = 30$ can be reached at different values of transverse momentum p_T which is corresponding to different values of collision energy $\sqrt{s}$. The measurement of J/ψ inclusive cross section at Tevatron energies over the range $p_T > 20$ GeV/c could give information on asymptotic behavior of $\psi(z)$, verify the predictions and establish features of particle formation mechanism at high- p_T .

4. CONCLUSIONS

Experimental data on inclusive cross section for prompt J/ψ production in $\bar{p}p$ collisions at $\sqrt{s} = 1800$ and 1960 GeV obtained at Run I and Run II with the CDF detector at Fermilab were analyzed in the framework of z -presentation. Using the properties of z -scaling, the energy independence and F -dependence of z -presentations, the scaling function of J/ψ constructed over a wide z -range.

In z -presentation the scaling function $\psi(z)$ and scaling variable z are expressed via the experimental quantities, the invariant inclusive cross section $Ed^3\sigma/dp^3$ and the multiplicity density of charged particles $\rho(s, \eta)$.

The physical interpretation of the scaling function ψ as a probability density to produce a particle with the formation length z is given. The quantity z has the property of the fractal measure, and $\delta_{1,2}$ is the fractal dimension describing the intrinsic structure of the interacting constituents revealed at high energies.

The prompt J/ψ spectra over a wide p_T and $\sqrt{s}$ range are predicted. The kinematic regions preferable to search for z -scaling violation in $\bar{p}p$ collisions at $\sqrt{s} = 1960$ GeV are established.

Thus the results of our analysis confirm the concept of z -scaling, the possibility of its applicability for analysis of experimental high- p_T data and search for new physics phenomena in Run II at the Tevatron.

Acknowledgments

The author would like to thank Yu.Gotra for useful and stimulating discussions of CDF experimental data and obtained results.

-
- ¹ R.P. Feynman, Phys. Rev. Lett. **23**, 1415 (1969).
 - ² J.D. Bjorken, Phys. Rev. **179**, 1547 (1969); J.D. Bjorken, and E.A. Paschos, Phys. Rev. **185**, 1975 (1969).
 - ³ P. Bosted et al., Phys. Rev. Lett. **49**, 1380 (1972).
 - ⁴ J. Benecke et al., Phys. Rev. **188**, 2159 (1969).
 - ⁵ A.M. Baldin, Sov. J. Part. Nucl. **8**, 429 (1977).
 - ⁶ V.S. Stavinsky, Sov. J. Part. Nucl. **10**, 949 (1979).
 - ⁷ G.A. Leksin: Report No. ITEP-147, 1976; G.A. Leksin: in *Proceedings of the XVIII International Conference on High Energy Physics*, Tbilisi, Georgia, 1976, edited by N.N. Bogolubov *et al.*, (JINR Report No. D1,2-10400, Tbilisi, 1977), p. A6-3.
 - ⁸ Z. Koba, H.B. Nielsen, and P. Olesen, Nucl. Phys. **B 40**, 317 (1972).
 - ⁹ V.A. Matveev, R.M. Muradyan, and A.N. Tavkhelidze, Part. Nuclei **2** 7 (1971); Lett. Nuovo Cimento **5**, 907 (1972); Lett. Nuovo Cimento **7**, 719 (1973).
 - ¹⁰ S. Brodsky, and G. Farrar, Phys. Rev. Lett. **31**, 1153 (1973); Phys. Rev. **D 11**, 1309 (1975).
 - ¹¹ I.Zborovský, Yu.A.Panebratsev, M.V.Tokarev, and G.P.Škoro, Phys. Rev. **D 54**, 5548 (1996); I.Zborovský, M.V.Tokarev, Yu.A.Panebratsev, and G.P.Škoro, Phys. Rev. **C59**, 2227 (1999); M.V.Tokarev, T.G.Dedovich, Int. J. Mod. Phys. **A15**, 3495 (2000); M.V.Tokarev, O.V.Rogachevski, T.G.Dedovich, J. Phys. G: Nucl. Part. Phys. **26**, 1671 (2000); M.V.Tokarev, O.V.Rogachevski, and T.G.Dedovich, Preprint No. E2-2000-90, JINR (Dubna, 2000); M.Tokarev, I.Zborovský, Yu.Panebratsev, and G.Skoro, Int. J. Mod. Phys. **A16**, 1281 (2001); M.Tokarev, hep-ph/0111202; M.Tokarev, D.Toivonen, hep-ph/0209069; G.P.Skoro, M.V.Tokarev, Yu.A.Panebratsev, and I.Zborovský, hep-ph/0209071; M.V.Tokarev, G.L.Efimov, and D.E.Toivonen, Physics of Atomic Nuclei, **67**, 564 (2004).
 - ¹² M.V.Tokarev, JINR Preprint E2-98-92, Dubna, 1998; JINR Preprint E2-98-161, Dubna, 1998; M.V.Tokarev, E.V.Potrebenikova, Computer Physics Communications **117**, 229 (1999); M.Tokarev, G.Efimov, hep-ph/0209013.
 - ¹³ F.Abe et al., Phys. Rev. Lett. **79**, 572 (1997); **69**, 3704 (1992); **79**, 578 (1997).
 - ¹⁴ Yu.Gotra: Ph.D. Thesis (Joint Institute for Nuclear Research, Dubna, Russia, 2004).
 - ¹⁵ R.Baier, and R.Ruckl, Z. Phys. **C19**, 251 (1983).
 - ¹⁶ E.Braaten, and S.Fleming, Phys. Rev. Lett. **74**, 3327 (1995); P.Cho, and M.Wise, Phys. Lett. **B346**, 129 (1995).
 - ¹⁷ M.Cacciari et al., Phys. Lett. **B356**, 553 (1995).
 - ¹⁸ P.Cho, and A.K.Leibovich, Phys. Rev. **D53**, 6203 (1996).
 - ¹⁹ M.Banner et al., Phys. Lett. **B115**, 59 (1982).




FIG. 1: (a) The scaling function of π^0 - and prompt J/ψ -mesons produced in $\bar{p}p$ collisions in the central rapidity range at $\sqrt{s} = 540$ and $1800, 1960$ GeV, respectively. Experimental data on inclusive cross section are taken from^{13,14,19}. (b) The inclusive differential cross sections of prompt J/ψ -mesons produced in $\bar{p}p$ collisions at $\sqrt{s} = 63, 200, 630, 1800$ and 1960 GeV in the central rapidity range as a functions of the transverse momentum p_T . Points are experimental data on inclusive cross section taken from^{13,14}. Dashed lines are predicted results based on z -scaling.




FIG. 2: The dependence of the scaling variable z on the transverse momentum p_T of J/ψ production in $\bar{p}p$ and pp collisions at $\sqrt{s} = 63 - 14000$ GeV and the angle $\theta_{\text{cms}} = 90^\circ$.

z -Scaling and Prompt J/ψ Production at High- p_T in $\bar{p}p$ at Tevatron

M.V. Tokarev*

Joint Institute for Nuclear Research, Dubna, Moscow region, 141980, Russia

New data on high- p_T spectra of J/ψ measured by the CDF Collaboration at the Tevatron are analyzed in the framework of z -scaling. F -dependence of z -presentation of the data is used to construct the scaling function of J/ψ . The energy independence of the scaling function is used to predict inclusive cross section of prompt J/ψ production in $\bar{p}p$ collisions at high- p_T . The concept of z -scaling is considered to reflect the general features of high- p_T particle production in hadron-hadron and hadron-nucleus collisions. Violation of z -scaling is suggested to consider as a signature of new physics.

1. INTRODUCTION

Primary goal of all new accelerators is to discover new physics phenomena. They can give impulse for development of physics theory and extend our understanding of micro world. The origin of particle structure at small scales, distributions of mass, charge and spin in space-time, features of particle-to-particle transitions, are of the most enigmatic and least understandable physics phenomena. Therefore, search for scaling regularities in high energy particle collisions is a subject of intense investigations¹⁻¹⁰.

Properties of particle formation are believed to reveal themselves more clearly at high energy $\sqrt{s}$ and transverse momenta p_T . It is considered also that partons produced in hard scattering retain information about primary collision during hadronization. It is known that the mechanism of particle formation is modified by nuclear medium and can be sensitive to the phase transition of nuclear matter. Therefore, the features of high- p_T single inclusive particle spectra of hadron-hadron and hadron-nucleus collisions are of interest to search for new physics phenomena in elementary processes (quark compositeness, fractal space-time, extra dimensions), signatures of exotic state of nuclear matter (phase transitions, quark-gluon plasma), complementary restrictions for theory (nonperturbative QCD effects, physics beyond Standard Model).

New feature (z -scaling) of high- p_T particle production in hadron-hadron and hadron-nucleus collisions at high energies was established in^{11,12}. The scaling function ψ and scaling variable z are expressed via experimental quantities such as the inclusive cross section $Ed^3\sigma/dp^3$ and the multiplicity density of charged particles $dN/d\eta$. The z -presentation of data is found to reveal symmetry properties (energy independence, A- and F-dependence, power law). The function $\psi(z)$ is interpreted as the probability density to produce a particle with a formation length z . The concept of z -scaling and the method of data analysis are developed for description of different particles (charged and neutral hadrons, direct photons jets) produced in high energy hadron-hadron and hadron-nucleus interactions. The proposed method is complementary to a method of direct calculations developed in the framework of QCD and methods based on the Monte Carlo generators. Therefore we consider that the use of the method allow to reduce some theoretical uncertainties which are ambiguously estimated by theory.

Recently new data¹⁴ on cross section of prompt J/ψ production in $\bar{p}p$ collisions in the central rapidity range $|\eta_{J/\psi}| < 0.6$ at $\sqrt{s} = 1.96$ TeV are measured by the CDF Collaboration. The data and the other one¹³ obtained in Run I give new possibility to verify models of J/ψ production mechanisms. In the color singlet model¹⁵ the charmonium meson retains the quantum numbers of the produced $\bar{c}c$ pair and each J/ψ state is directly produced in hard scattering of color singlet via the corresponding subprocess. As shown in¹³ the model predictions are in a disagreement with experimental data on J/ψ and $\psi(2S)$ production rates. A color octet model^{16,17} was suggested to explain the discrepancy. The color octet mechanism takes into account the production of $\bar{c}c$ pair in a color octet configuration. The color octet state evolves into a color singlet state via emission of a soft gluon. The color octet contributions were found in¹⁸ from a fit to CDF data¹³.

We use a complementary method of data analysis and present the results of analysis of CDF data¹⁴ on prompt J/ψ production in $\bar{p}p$ collisions in Run I and Run II at Tevatron in the framework of z -scaling. Properties of z -scaling, the energy independence and F -dependence, are used to construct the scaling function for J/ψ . The inclusive cross sections of J/ψ production in $\bar{p}p$ collisions at high- p_T are estimated. The results can be useful to test phenomenological models of J/ψ production mechanism and obtain new restrictions on a gluon distribution function. The possibility to search for the scaling violation at the Tevatron is suggested as well.

* e-mail: tokarev@sunhe.jinr.ru

2. Z-SCALING

In the section underlying ideas of z -scaling, a general scheme of data z -presentation and physical meaning of the scaling function $\psi(z)$ and the scaling variable z are discussed.

2.1. Basic principles: locality, self-similarity, fractality

The idea of z -scaling is based on the assumptions⁶ that gross feature of inclusive particle distribution of the process (1) at high energies can be described in terms of the corresponding kinematic characteristics

$$M_1 + M_2 \rightarrow m_1 + X \quad (1)$$

of the constituent subprocess written in the symbolic form (2)

$$(x_1 M_1) + (x_2 M_2) \rightarrow m_1 + (x_1 M_1 + x_2 M_2 + m_2) \quad (2)$$

satisfying the condition

$$(x_1 P_1 + x_2 P_2 - p)^2 = (x_1 M_1 + x_2 M_2 + m_2)^2. \quad (3)$$

The equation is the expression of locality of hadron interaction at constituent level. The x_1 and x_2 are fractions of the incoming momenta P_1 and P_2 of the colliding objects with the masses M_1 and M_2 . They determine the minimum energy, which is necessary for production of the secondary particle with the mass m_1 and the four-momentum p . The parameter m_2 is introduced to satisfy the internal conservation laws (for baryon number, isospin, strangeness, and so on).

The equation (3) reflects minimum recoil mass hypothesis in the elementary subprocess. To connect kinematic and structural characteristics of the interaction, the quantity Ω is introduced. It is chosen in the form

$$\Omega(x_1, x_2) = m(1 - x_1)^{\delta_1}(1 - x_2)^{\delta_2}, \quad (4)$$

where m is a mass constant and δ_1 and δ_2 are factors relating to the anomalous fractal dimensions of the colliding objects. The fractions x_1 and x_2 are determined to maximize the value of $\Omega(x_1, x_2)$, simultaneously fulfilling the condition (3)

$$d\Omega(x_1, x_2)/dx_1|_{x_2=x_2(x_1)} = 0. \quad (5)$$

The fractions x_1 and x_2 are equal to unity along the phase space limit and cover the full phase space accessible at any energy.

Self-similarity is a scale-invariant property connected with dropping of certain dimensional quantities out of physical picture of the interactions. It means that dimensionless quantities for the description of physical processes are used. The scaling function $\psi(z)$ depends in a self-similar manner on the single dimensionless variable z . It is expressed via the invariant cross section $E d^3\sigma/dp^3$ as follows

$$\psi(z) = -\frac{\pi s}{(dN/d\eta)\sigma_{in}} J^{-1} E \frac{d^3\sigma}{dp^3} \quad (6)$$

Here, s is the center-of-mass collision energy squared, σ_{in} is the inelastic cross section, J is the corresponding Jacobian. The factor J is the known function of the kinematic variables, the momenta and masses of the colliding and produced particles.

The function $\psi(z)$ is normalized as follows

$$\int_0^\infty \psi(z) dz = 1. \quad (7)$$

The relation allows us to interpret the function $\psi(z)$ as a probability density to produce a particle with the corresponding value of the variable z . We would like to emphasize that the existence of the function $\psi(z)$ depending on a single dimensionless variable z and revealing scaling properties is not evident in advance. Therefore the method proposed for data analysis in z -presentation could be only proved a posteriori.

Principle of fractality states that variables used in the description of the process diverge in terms of the resolution. This property is characteristic for the scaling variable

$$z = z_0 \Omega^{-1}, \quad (8)$$

where

$$z_0 = \sqrt{\hat{s}_\perp}/(dN/d\eta). \quad (9)$$

The variable z has character of a fractal measure. For the given production process (1), its finite part z_0 is the ratio of the transverse energy released in the binary collision of constituents (2) and the average multiplicity density $dN/d\eta|_{\eta=0}$. The divergent part Ω^{-1} describes the resolution at which the collision of the constituents can be singled out of this process. The $\Omega(x_1, x_2)$ represents relative number of all initial configurations containing the constituents which carry fractions x_1 and x_2 of the incoming momenta. The δ_1 and δ_2 are the anomalous fractal dimensions of the colliding objects (hadrons or nuclei). The momentum fractions x_1 and x_2 are determined in a way to minimize the resolution $\Omega^{-1}(x_1, x_2)$ of the fractal measure z with respect to all possible sub-processes (2) subjected to the condition (3). The variable z was interpreted as a particle formation length.

The scaling function of high- p_T particle production, as shown below, is described by the power law, $\psi(z) \sim z^{-\beta}$. Both quantities, ψ and z , are scale dependent. Therefore we consider the high energy hadron-hadron interactions as interactions of fractals. In the asymptotic region the internal structure of particles, interactions of their constituents and mechanism of real particle formation manifest self-similarity and fractality over a wide scale range.

2.2. Prompt J/ψ production in $\bar{p}p$

In this section we analyze the data^{13,14} on inclusive cross section of prompt J/ψ production in $\bar{p}p$ collisions. Firstly the experimental data¹⁹ on inclusive cross section of π^0 -mesons produced in $\bar{p}p$ collisions obtained by the UA1 Collaboration at $\sqrt{s} = 540$ GeV are used to construct z -presentation of the data. The scaling function of π^0 is shown in Fig.1(a). The dashed line is the fit of the data. As seen from Fig.1(a) the scaling function is described by the power law, $\psi(z) \sim z^{-\beta}$, on the log-log scale. The value of the slope parameter β is found to be 5.77 ± 0.02 over a wide p_T range. Secondly the F -dependence of z -scaling is used to compare the scaling function of π^0 and J/ψ . We assume that scaling functions of produced particles with different flavor content has the same asymptotic behavior^{11,12}. The transformation of variable z and scaling function ψ in the form

$$z \rightarrow \alpha_F \cdot z, \quad \psi \rightarrow \alpha_F^{-1} \cdot \psi \quad (10)$$

was used. The transformation parameter α_F is found to be 2.25. As seen from Fig.1(a) the scaling function of J/ψ at $z > 10$ reveals the asymptotic behavior such as the similar one for π^0 . We would like to note that the CDF Run I and Run II data demonstrate a good matching in the overlapping region in z -presentation. The power law for the scaling function is observed up to $z \simeq 40$. Therefore the kinematic range of more preferable for search for the scaling violation is found to be for the range $p_T > 30$ GeV/c at $\sqrt{s} = 1960$ GeV.

The energy independence of the constructed scaling function is used to calculate the dependence of inclusive cross section $Ed^3\sigma/dp^3$ of J/ψ production in $\bar{p}p$ collisions on the transverse momentum p_T in the central rapidity range, $\theta \simeq 90^\circ$, at the energy $\sqrt{s} = 63, 200, 630, 1800$ and 1960 GeV. The calculated results are shown by the dashed lines in Fig.1(b). Points ($\star, +$) shown in Fig.1(b) are the corresponding CDF Run I and Run II data. As seen from Fig.1(b) the strong dependence of inclusive cross section on $\sqrt{s}$ increases with p_T .

3. $z - p_T$ PLOT

The scaling function $\psi(z)$ demonstrates the universality over a wide kinematic range. It describes two regimes of particle formation observed at higher energies. The first one reveals exponential behavior at low z and the second one is described by the power law, $\psi(z) \sim z^{-\beta}$, at high z .

Asymptotic behavior of the scaling function is described by the slope parameter β , which depends on the anomalous fractal dimensions $\delta_{1,2}$ of colliding objects. We assume that new physics effects at small scales (high- z) should be characterized by the change of the fractal dimension (" δ -jump") and consequently the other value of the slope parameter β at high z . To determine the kinematic range which is of more preferable for experimental searching for z -scaling violation the $z - p_T$ plot can be used. Figure 2 shows the $z - p_T$ plot, the dependence of the variable z on the transverse momentum p_T of J/ψ -mesons produced in $\bar{p}p$ collisions at $\sqrt{s} = 63 - 14000$ GeV and $\theta_{cms} = 90^\circ$. As seen from Fig.1(a) the asymptotic value of the scaling function $\psi(z)$ at $z = 30$ can be reached at different values of transverse momentum p_T which is corresponding to different values of collision energy $\sqrt{s}$. The measurement of J/ψ inclusive cross section at Tevatron energies over the range $p_T > 20$ GeV/c could give information on asymptotic behavior of $\psi(z)$, verify the predictions and establish features of particle formation mechanism at high- p_T .

4. CONCLUSIONS

Experimental data on inclusive cross section for prompt J/ψ production in $\bar{p}p$ collisions at $\sqrt{s} = 1800$ and 1960 GeV obtained at Run I and Run II with the CDF detector at Fermilab were analyzed in the framework of z -presentation. Using the properties of z -scaling, the energy independence and F -dependence of z -presentations, the scaling function of J/ψ constructed over a wide z -range.

In z -presentation the scaling function $\psi(z)$ and scaling variable z are expressed via the experimental quantities, the invariant inclusive cross section $Ed^3\sigma/dp^3$ and the multiplicity density of charged particles $\rho(s, \eta)$.

The physical interpretation of the scaling function ψ as a probability density to produce a particle with the formation length z is given. The quantity z has the property of the fractal measure, and $\delta_{1,2}$ is the fractal dimension describing the intrinsic structure of the interacting constituents revealed at high energies.

The prompt J/ψ spectra over a wide p_T and $\sqrt{s}$ range are predicted. The kinematic regions preferable to search for z -scaling violation in $\bar{p}p$ collisions at $\sqrt{s} = 1960$ GeV are established.

Thus the results of our analysis confirm the concept of z -scaling, the possibility of its applicability for analysis of experimental high- p_T data and search for new physics phenomena in Run II at the Tevatron.

Acknowledgments

The author would like to thank Yu.Gotra for useful and stimulating discussions of CDF experimental data and obtained results.

-
- ¹ R.P. Feynman, Phys. Rev. Lett. **23**, 1415 (1969).
 - ² J.D. Bjorken, Phys. Rev. **179**, 1547 (1969); J.D. Bjorken, and E.A. Paschos, Phys. Rev. **185**, 1975 (1969).
 - ³ P. Bosted et al., Phys. Rev. Lett. **49**, 1380 (1972).
 - ⁴ J. Benecke et al., Phys. Rev. **188**, 2159 (1969).
 - ⁵ A.M. Baldin, Sov. J. Part. Nucl. **8**, 429 (1977).
 - ⁶ V.S. Stavinsky, Sov. J. Part. Nucl. **10**, 949 (1979).
 - ⁷ G.A. Leksin: Report No. ITEF-147, 1976; G.A. Leksin: in *Proceedings of the XVIII International Conference on High Energy Physics*, Tbilisi, Georgia, 1976, edited by N.N. Bogolubov *et al.*, (JINR Report No. D1,2-10400, Tbilisi, 1977), p. A6-3.
 - ⁸ Z. Koba, H.B. Nielsen, and P. Olesen, Nucl. Phys. **B 40**, 317 (1972).
 - ⁹ V.A. Matveev, R.M. Muradyan, and A.N. Tavkhelidze, Part. Nuclei **2** 7 (1971); Lett. Nuovo Cimento **5**, 907 (1972); Lett. Nuovo Cimento **7**, 719 (1973).
 - ¹⁰ S. Brodsky, and G. Farrar, Phys. Rev. Lett. **31**, 1153 (1973); Phys. Rev. **D 11**, 1309 (1975).
 - ¹¹ I.Zborovský, Yu.A.Panebratsev, M.V.Tokarev, and G.P.Škoro, Phys. Rev. **D 54**, 5548 (1996); I.Zborovský, M.V.Tokarev, Yu.A.Panebratsev, and G.P.Škoro, Phys. Rev. **C59**, 2227 (1999); M.V.Tokarev, T.G.Dedovich, Int. J. Mod. Phys. **A15**, 3495 (2000); M.V.Tokarev, O.V.Rogachevski, T.G.Dedovich, J. Phys. G: Nucl. Part. Phys. **26**, 1671 (2000); M.V.Tokarev, O.V.Rogachevski, and T.G.Dedovich, Preprint No. E2-2000-90, JINR (Dubna, 2000); M.Tokarev, I.Zborovský, Yu.Panebratsev, and G.Skoro, Int. J. Mod. Phys. **A16**, 1281 (2001); M.Tokarev, hep-ph/0111202; M.Tokarev, D.Toivonen, hep-ph/0209069; G.P.Skoro, M.V.Tokarev, Yu.A.Panebratsev, and I.Zborovský, hep-ph/0209071; M.V.Tokarev, G.L.Efimov, and D.E.Toivonen, Physics of Atomic Nuclei, **67**, 564 (2004).
 - ¹² M.V.Tokarev, JINR Preprint E2-98-92, Dubna, 1998; JINR Preprint E2-98-161, Dubna, 1998; M.V.Tokarev, E.V.Potrebenikova, Computer Physics Communications **117**, 229 (1999); M.Tokarev, G.Efimov, hep-ph/0209013.
 - ¹³ F.Abe et al., Phys. Rev. Lett. **79**, 572 (1997); **69**, 3704 (1992); **79**, 578 (1997).
 - ¹⁴ Yu.Gotra: Ph.D. Thesis (Joint Institute for Nuclear Research, Dubna, Russia, 2004).
 - ¹⁵ R.Baier, and R.Ruckl, Z. Phys. **C19**, 251 (1983).
 - ¹⁶ E.Braaten, and S.Fleming, Phys. Rev. Lett. **74**, 3327 (1995); P.Cho, and M.Wise, Phys. Lett. **B346**, 129 (1995).
 - ¹⁷ M.Cacciari et al., Phys. Lett. **B356**, 553 (1995).
 - ¹⁸ P.Cho, and A.K.Leibovich, Phys. Rev. **D53**, 6203 (1996).
 - ¹⁹ M.Banner et al., Phys. Lett. **B115**, 59 (1982).




FIG. 1: (a) The scaling function of π^0 - and prompt J/ψ -mesons produced in $\bar{p}p$ collisions in the central rapidity range at $\sqrt{s} = 540$ and $1800, 1960$ GeV, respectively. Experimental data on inclusive cross section are taken from^{13,14,19}. (b) The inclusive differential cross sections of prompt J/ψ -mesons produced in $\bar{p}p$ collisions at $\sqrt{s} = 63, 200, 630, 1800$ and 1960 GeV in the central rapidity range as a functions of the transverse momentum p_T . Points are experimental data on inclusive cross section taken from^{13,14}. Dashed lines are predicted results based on z -scaling.




FIG. 2: The dependence of the scaling variable z on the transverse momentum p_T of J/ψ production in $\bar{p}p$ and pp collisions at $\sqrt{s} = 63 - 14000$ GeV and the angle $\theta_{\text{cms}} = 90^\circ$.